

Final approval of the article: HB

Statistical analysis: HB

Obtained funding: HB, WS

Overall responsibility: HB

## REFERENCES

- Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG, et al. Inter-society consensus for the management of peripheral arterial disease (TASC II). *J Vasc Surg* 2007;45(Suppl S):S5-67.
- Davies M. Critical limb ischemia: epidemiology. *Methodist DeBakey Cardiovasc J* 2012;8:10-4.
- Albers M, Romiti M, Brochado-Neto FC, De Luccia N, Pereira CA. Meta-analysis of popliteal-to-distal vein bypass grafts for critical ischemia. *J Vasc Surg* 2006;43:498-503.
- Kudo T, Chandra FA, Kwun WH, Haas BT, Ahn SS. Changing pattern of surgical revascularization for critical limb ischemia over 12 years: endovascular vs. open bypass surgery. *J Vasc Surg* 2006;44:304-13.
- Botti CF Jr, Ansel GM, Silver MJ, Barker BJ, South S. Percutaneous retrograde tibial access in limb salvage. *J Endovasc Ther* 2003;10:614-8.
- Spinosa DJ, Leung DA, Harthun NL, Cage DL, Fritz Angle J, Hagspiel KD, et al. Simultaneous antegrade and retrograde access for subintimal recanalization of peripheral arterial occlusion. *J Vasc Interv Radiol* 2003;14:1449-54.
- Manzi M, Palena LM. Retrograde percutaneous transmetatarsal artery access: new approach for extreme revascularization in challenging cases of critical limb ischemia. *Cardiovasc Interv Radiol* 2013;36:554-7.
- Jaffan AA, Chu V, Murphy TP. Fate of retrograde pedal access after subintimal arterial flossing with antegrade-retrograde intervention (SA-FARI) procedure in patients with intermittent claudication: preliminary experience in three patients. *J Vasc Interv Radiol* 2013;24:139-42.
- El-Sayed HF. Retrograde pedal/tibial artery access for treatment of infragenicular arterial occlusive disease. *Methodist DeBakey Cardiovasc J* 2013;9:73-8.
- Chang JC, Lin LS, Chiu CH. Static contrast technique for creating transpedal arterial access in patients with tibioperoneal occlusions. *J Vasc Surg* 2013;58:1120-2.
- Shimada Y, Kino N, Yano K, Tonomura D, Takehara K, Furubayashi K, et al. Transcollateral retrograde approach with rendezvous technique for recanalization of chronically occluded tibial arteries. *J Endovasc Ther* 2012;19:620-6.
- Schmidt A, Bausback Y, Piorkowski M, Werner M, Braunlich S, Ulrich M, et al. Retrograde recanalization technique for use after failed antegrade angioplasty in chronic femoral artery occlusions. *J Endovasc Ther* 2012;19:23-9.
- Ruzsa Z, Pinter L, Kolvenbach R. Retrograde transpedal stenting of the tibioperoneal trunk in critical limb ischemia. *Catheter Cardiovasc Interv* 2012;80:1105-11.
- Londono JC, Singh V, Martinez CA. Posterior tibial artery access using transradial techniques: Retrograde approach to inaccessible lower extremity lesions. *Catheter Cardiovasc Interv* 2012;79:1194-8.
- Alqahtani S, Kandeel AY, Rolf T, Frederic G, Qanadli SD. Case report: an unusual combined retrograde and antegrade transpedal subintimal recanalization of the infraginal arteries. *J Vasc Interv Radiol* 2012;23:1325-9.
- Hansen A, Krawczynski H, Lacher F. Retrograde transpedale recanalisation of tibial artery occlusion. *Vasa* 2009;38:249-53.
- Kawarada O, Yokoi Y, Sekii H, Higashiue S. Retrograde crossing through the pedal arch for totally occluded tibial artery. *J Intervent Cardiol* 2008;21:342-6.
- Fusaro M, Tashani A, Mollicelli N, Medda M, Inglese L, Biondi-Zoccai GG. Retrograde pedal artery access for below-the-knee percutaneous revascularisation. *J Cardiovasc Med* 2007;8:216-8.
- Downer J, Uderoi R. Percutaneous retrograde tibial access in the endovascular treatment of acute limb ischaemia: a case report. *Eur J Cardiovasc Surg* 2007;34:350-2.
- Ye M, Zhang H, Huang XZ, Zhang L, Shi YX, Liang W, et al. [Retrograde tibial approach for endovascular recanalization of popliteal and tibioperoneal artery occlusion]. *Zhonghua Wai Ke Za Zhi* 2013;51:710-4.
- Ruzsa Z, Nemes B, Bansaghi Z, Toth K, Kuti F, Kudrnova S, et al. Transpedal access after failed antegrade recanalization of complex below-the-knee and femoropopliteal occlusions in critical limb ischemia [published online ahead of print Oct 19, 2013]. *Catheter Cardiovasc Interv* doi: 10.1002/ccd.25262.
- Mustapha JA, Saab F, McGoff T, Heaney C, Diaz-Sandoval L, Sevenma M, et al. Tibio-pedal arterial minimally invasive retrograde revascularization in patients with advanced peripheral vascular disease: the TAMI technique, original case series [published online ahead of print Oct 6, 2013]. *Catheter Cardiovasc Interv* doi: 10.1002/ccd.25227.
- Palena LM, Manzi M. Extreme below-the-knee interventions: Retrograde transmetatarsal or transplanter arch access for foot salvage in challenging cases of critical limb ischemia. *J Endovasc Ther* 2012;19:805-11.
- Montero-Baker M, Schmidt A, Braunlich S, Ulrich M, Thieme M, Biamino G, et al. Retrograde approach for complex popliteal and tibioperoneal occlusions. *J Endovasc Ther* 2008;15:594-604.
- Fernandez N, McEnaney R, Marone LK, Rhee RY, Leers S, Makaroun M, et al. Multilevel versus isolated endovascular tibial interventions for critical limb ischemia. *J Vasc Surg* 2011;54:722-9.
- Shin SH, Baril D, Chaer R, Rhee R, Makaroun M, Marone L. Limitations of the outback ltd re-entry device in femoropopliteal chronic total occlusions. *J Vasc Surg* 2011;53:1260-4.
- Srivatsa SS, Edwards WD, Boos CM, Grill DE, Sangiorgi GM, Garratt KN, et al. Histologic correlates of angiographic chronic total coronary artery occlusions: influence of occlusion duration on neo-vascular channel patterns and intimal plaque composition. *J Am Coll Cardiol* 1997;29:955-63.
- Saito S. Different strategies of retrograde approach in coronary angioplasty for chronic total occlusion. *Catheter Cardiovasc Interv* 2008;71:8-19.
- Bazan HA. Think of the angiosome concept when revascularizing the patient with critical limb ischemia. *Catheter Cardiovasc Interv* 2010;75:837.
- Iida O, Nanto S, Uematsu M, Ikeoka K, Okamoto S, Dohi T, et al. Importance of the angiosome concept for endovascular therapy in patients with critical limb ischemia. *Catheter Cardiovasc Interv* 2010;75:830-6.
- Zhuang KD, Tan SG, Tay KH. The "safari" technique using retrograde access via peroneal artery access. *Cardiovasc Interv Radiol* 2012;35:927-31.
- Feiring AJ, Krahn M, Nelson L, Wesolowski A, Eastwood D, Szabo A. Preventing leg amputations in critical limb ischemia with below-the-knee drug-eluting stents: the PARADISE (preventing amputations using drug eluting stents) trial. *J Am Coll Cardiol* 2010;55:1580-9.
- Gasper WJ, Runge SJ, Owens CD. Management of infrapopliteal peripheral arterial occlusive disease. *Curr Treat Options Cardiovasc Med* 2012;14:136-48.

Submitted Dec 17, 2013; accepted Feb 13, 2014.

## DISCUSSION

**Dr Cynthia K. Shortell** (*Durham, NC*). Congratulations to the authors on this excellent paper about a challenging clinical and technical problem; namely critical limb ischemia due to tibial disease in patients who are poor surgical risk.

As vascular surgeons, we are increasingly called upon to achieve limb salvage in more and more difficult situations, be it unfavorable patient anatomy, patient physiology, or patient physique, and we need ever-greater numbers of tools in our toolbox in order

to accomplish this. This paper describes the Ochsner group's experience with just such a tool; namely, retrograde pedal access for patients with tibial artery occlusive disease. While this technique has been described previously, midterm results have not been reported previously. The authors describe the use of this technique in patients in whom a standard antegrade approach had been tried and had failed, and report favorable technical and clinical success rates.

I thoroughly enjoyed reading this well-written manuscript and appreciate receiving it from the authors well in advance of the meeting. I would also like to thank the authors for providing me with a copy of the manuscript in a very timely manner. I have several questions for the authors:

First, can you be more specific about the unsuitability for surgery? Were there medical comorbidities or patient-related issues such as obesity? How was this determined? Were there specific criteria or was it surgeon-specific?

Your manuscript suggests that all patients had an initial attempt at traditional antegrade access. Was this indeed the case, and if so, was the pedal access attempted at the same or a later setting?

In four patients, the procedure was technically unsuccessful. What happened to these patients?

You had a limb salvage rate of 77% at 17 months. Did you assess patency of the revascularization, and if so, what was it?

Given the results of this study will you change your practice to offer pedal access as the initial procedure in some cases now?

**Dr Hernan Bazan.** Thank you, Dr Shortell, for the nice summary and the very insightful questions. The first question you asked is about the unsuitability of the patients. We did not do any standardized risk stratification. Patient suitability was left to one of the three board-certified vascular surgeons who deemed whether or not the patient was a bypass candidate. Patient factors,

such as ischemic cardiomyopathy with a severely depressed ejection fraction or dialysis-dependence with a poor functional status, are some examples. In terms of whether or not we brought the patients back for a retrograde pedal access during the initial angiogram, I can only recall one or two instances where the patient was brought back after a failed antegrade access. This does bring up another point, which we were discussing earlier: in these cases that involve a retrograde pedal access after a failed antegrade approach, what we have found is that after one attempt to revascularized via an antegrade approach for an hour or one hour and a half, the techs in the endovascular suite are then told "okay, let's prep the foot," and everybody in the endo suite has to be prepared to work an additional hour or so. Hence, it certainly does add more time, but I can recall one or two patients where they were brought back. These were primarily cases in which we were trying to avoid a large contrast exposure in patients with compromised renal function.

You also asked a question about the four patients we were not able to revascularized; there was major limb loss in three of them. The fourth patient had rest pain, and he subsequently underwent a successful revascularization through a pedal bypass. He was the only one out of the 13 that had a bypass, as none of the other patients were bypass candidates.

The last question you asked: Would this change our practice? I think, as it was mentioned earlier and mentioned on the first day of our conference, we always teach our trainees to individualize treatment to the patient. I believe we should optimize our attempt to do a tibial bypass when we can because it is a great operation, very durable, and to date, nothing else has similar long-term patency. Although we do in fact attempt an endovascular approach first in the majority of patients, there are some who are clearly good bypass candidates and no endoluminal attempt at revascularization is made; they go on to a tibial bypass.